



*The Medical **Bulletin***

In Cardiology

Specifically, patients with uncomplicated pericarditis have a “pleuritic” type of chest pain, sometimes radiating to the one or both shoulders that decreases in intensity when the patient sits up and leans forward. Occasionally, however, the patient will describe a dull, oppressive discomfort much like that of angina or myocardial infarction. In this setting, listen carefully for a pericardial friction rub, specific for acute pericarditis. On the electrocardiogram, look for widespread ST elevation or P-R depression

Dr. Baktavatchalam

In Cardiology

CARDIOLOGY QUESTIONS

1. Anti arrhythmic drug causing lupus like syndrome
2. Drug used to suppress frequent ventricular tachycardia in brugada syndrome
3. Cardiac lesion seen in ankylosing spondylitis
4. Ionotropic agent does not have ionotropic effect
5. Complications are least in radiofrequency ablation of which accessory pathway?
6. Myocardial bridging is most common in which coronary artery
7. Most common game associated with commotiocordis
8. Cabrera sigh is seen in
9. CAD patient on treatment develops skin rash, leucopenia, thrombocytopenia-drug responsible?
10. Most potent endogenous cardiostimulatory peptide?

CARDIOLOGY ANSWERS

1. Procainamide
2. Quinidine
3. Aortic regurgitation
4. Nesiritide
5. Left lateral pathway
6. LAD
7. Baseball
8. Old MI +LBBB
9. Ticlopidine
10. Urotensin